

REMARKS ON STRUCTURE OF CAT(0) GROUPS

TETSUYA HOSAKA

ABSTRACT. In this paper, we investigate finitely generated groups of isometries of CAT(0) spaces containing some central hyperbolic isometry, and we provide some remarks on structure of CAT(0) groups.

1. INTRODUCTION AND PRELIMINARIES

The purpose of this paper is to investigate finitely generated groups of isometries of CAT(0) spaces and CAT(0) groups. Definition and detail of CAT(0) spaces are found in [3] and [9].

Let X be a metric space and let γ be an isometry of X . Then the *translation length* of γ is defined as $|\gamma| = \inf\{d(x, \gamma x) \mid x \in X\}$, and the *minimal set* of γ is defined as $\text{Min}(\gamma) = \{x \in X \mid d(x, \gamma x) = |\gamma|\}$. An isometry γ is said to be *semi-simple* if $\text{Min}(\gamma)$ is non-empty. Also an isometry γ is called

- (1) *elliptic* if γ has a fixed point,
- (2) *hyperbolic* if γ is semi-simple and not elliptic, and
- (3) *parabolic* if γ is not semi-simple.

(cf. [3, Chapter II.6]).

We first show the following theorem by observing the proof of [3, Theorem II.6.12] in Section 2.

Theorem 1.1. *Let X be a CAT(0) space and let Γ be a finitely generated group acting by isometries on X . If the center of Γ contains a hyperbolic isometry γ_0 of X , then there exist a normal subgroup $\Gamma' \subset \Gamma$, an element $\delta_0 \in \Gamma$ and a number $k_0 \in \mathbb{N}$ such that*

- (i) $\Gamma = \Gamma' \rtimes \langle \delta_0 \rangle$,
- (ii) $\Gamma' \rtimes \langle \delta_0^{k_0} \rangle = \Gamma' \times \langle \gamma_0 \rangle$ is a finite-index subgroup of Γ and
- (iii) Γ/Γ' is isomorphic to $\mathbb{Z}$.

Date: April 25, 2010.

2000 Mathematics Subject Classification. 20F65; 57M07.

Key words and phrases. CAT(0) group; CAT(0) space; isometry of CAT(0) space.

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 21740037).

A *geometric* action on a CAT(0) space is an action by isometries which is proper ([3, p.131]) and cocompact. A group Γ is called a *CAT(0) group*, if Γ acts geometrically on some CAT(0) space. We note that every CAT(0) space on which some group acts geometrically is a proper space ([3, p.132]). Also we note that CAT(0) groups are finitely presented (cf. [3, Corollary I.8.11]).

For example, Bieberbach groups ([3, p.246], [4]), crystallographic groups ([4]), Coxeter groups and their torsion-free subgroups of finite-index ([6], [7], [18]) and fundamental groups of compact geodesic spaces of non-positive curvature ([3, p.159, p.237]) are CAT(0) groups. In particular, fundamental groups of Riemannian manifolds of non-positive sectional curvature are CAT(0) groups. Moreover, M. W. Davis [6] has constructed a closed aspherical manifold of dimension $n \geq 5$ whose universal covering is not homeomorphic to $\mathbb{R}^n$ ([6], [8]). The fundamental groups of these exotic manifolds are also CAT(0) groups.

On structure of CAT(0) groups, we obtain the following theorem from Theorem 1.1 in Section 2.

Theorem 1.2. *Let Γ be a CAT(0) group. Then there exist subgroups $\Gamma = \Gamma_0 \supset \Gamma_1 \supset \cdots \supset \Gamma_n$, elements $\delta_{i+1}, \gamma_{i+1} \in \Gamma_i$ and $k_{i+1} \in \mathbb{N}$ for $i = 0, \dots, n-1$ such that*

- (1) γ_{i+1} is an element of the center of Γ_i with the order $o(\gamma_{i+1}) = \infty$ for $i = 0, \dots, n-1$,
- (2) $\Gamma_i = \Gamma_{i+1} \rtimes \langle \delta_{i+1} \rangle$ for $i = 0, \dots, n-1$,
- (3) $\Gamma_{i+1} \rtimes \langle \delta_{i+1}^{k_{i+1}} \rangle = \Gamma_{i+1} \times \langle \gamma_{i+1} \rangle$ is a finite-index subgroup of Γ_i ,
- (4) Γ_i / Γ_{i+1} is isomorphic to $\mathbb{Z}$ for $i = 0, \dots, n-1$,
- (5) $\Gamma = (\cdots (((\Gamma_n \rtimes \langle \delta_n \rangle) \rtimes \langle \delta_{n-1} \rangle) \rtimes \langle \delta_{n-2} \rangle) \cdots) \rtimes \langle \delta_1 \rangle$,
- (6) Γ_n has finite center, and
- (7) $\Gamma_n \times A$ is a finite-index subgroup of Γ where $A = \langle \gamma_1 \rangle \times \cdots \times \langle \gamma_n \rangle$ which is isomorphic to $\mathbb{Z}^n$.

In Section 3, we provide some examples and remarks on CAT(0) groups.

2. ON FINITELY GENERATED GROUPS OF ISOMETRIES OF CAT(0) SPACES AND STRUCTURE OF CAT(0) GROUPS

We first note that we obtain the following lemma from the proof of [3, Theorem II.6.12].

Lemma 2.1 (cf. [3, Theorem II.6.12]). *Let X be a CAT(0) space and let Γ be a finitely generated group acting by isometries on X . If the center of Γ contains a hyperbolic isometry γ_0 of X , then there exists*

a homomorphism $\phi : \Gamma \rightarrow \mathbb{Z}$ such that $\phi(A)$ is non-trivial, where $A = \langle \gamma_0 \rangle \cong \mathbb{Z}$.

Here we show Theorem 1.1 from Lemma 2.1 by observing the proof of [3, Theorem II.6.12].

Proof of Theorem 1.1. Let X be a CAT(0) space and let Γ be a finitely generated group acting by isometries on X . Suppose that the center of Γ contains a hyperbolic isometry γ_0 of X . Then $A = \langle \gamma_0 \rangle \cong \mathbb{Z}$ is a central subgroup of Γ .

By Lemma 2.1, we obtain a homomorphism $\phi : \Gamma \rightarrow \mathbb{Z}$ such that $\phi(A)$ is non-trivial.

Here γ_0 generates A and $\phi(\gamma_0)$ generates $\phi(A)$. Let $H_0 = \phi^{-1}(\phi(A))$ which is a finite-index subgroup of Γ . Then as the proof of [3, Theorem II.6.12], the map $\phi(\gamma_0) \mapsto \gamma_0$ splits $\phi|_{H_0}$, hence we obtain that

$$H_0 = \ker \phi \times \langle \gamma_0 \rangle \quad (1),$$

since γ_0 is central.

Here also we note that $\phi(\Gamma)$ is a non-trivial subgroup of $\mathbb{Z}$. Let $\delta_0 \in \Gamma$ such that $\phi(\delta_0)$ generates $\phi(\Gamma)$. Then the map $\phi(\delta_0) \mapsto \delta_0$ splits ϕ , hence we obtain that

$$\Gamma = \ker \phi \rtimes \langle \delta_0 \rangle \quad (2),$$

since $\ker \phi$ is a normal subgroup of Γ .

Now $\phi(A)$ is a subgroup of $\phi(\Gamma)$ which is isomorphic to $\mathbb{Z}$ and $\phi(\delta_0)$ generates $\phi(\Gamma)$. Hence there exists a number $k_0 \in \mathbb{N}$ such that $\phi(\delta_0^{k_0})$ generates $\phi(A)$. Then $\phi^{-1}(\phi(\langle \delta_0^{k_0} \rangle)) = \phi^{-1}(\phi(A)) = H_0$ and the map $\phi(\delta_0^{k_0}) \mapsto \delta_0^{k_0}$ splits $\phi|_{H_0}$, hence we obtain that

$$H_0 = \ker \phi \rtimes \langle \delta_0^{k_0} \rangle \quad (3).$$

Let $\Gamma' = \ker \phi$. Then $\Gamma = \Gamma' \rtimes \langle \delta_0 \rangle$ by (2) and $H_0 = \Gamma' \rtimes \langle \gamma_0 \rangle = \Gamma' \rtimes \langle \delta_0^{k_0} \rangle$ is a finite-index subgroup of Γ by (1) and (3). Also $\Gamma/\Gamma' = \langle \Gamma'\delta_0 \rangle$ is isomorphic to $\mathbb{Z}$ by the construction. $\square$

The following results are known and we use these in proof of Theorem 1.2.

Lemma 2.2 ([3, Corollary I.8.11]). *A group is finitely presented if and only if it acts properly and cocompactly by isometries on a simply-connected geodesic space.*

Lemma 2.3 ([3, Proposition II.6.7]). *Let X be a complete CAT(0) space and let γ be an isometry of X . Then γ is elliptic if and only if γ has a bounded orbit.*

Lemma 2.4 ([3, Proposition II.6.10 (2)]). *If a group Γ acts properly and cocompactly by isometries on a metric space X , then every element of Γ is a semi-simple isometry of X .*

Lemma 2.5 ([3, Corollary II.7.6]). *Every abelian subgroup of a $CAT(0)$ group is finitely generated.*

The following theorem is called the Flat Torus Theorem.

Theorem 2.6 ([2, Proposition 1.1], [3, Theorem II.7.1]). *Let G be a group and let A be a free abelian group of rank n . Suppose that $\Gamma = G \times A$ acts geometrically on a $CAT(0)$ space X . Then*

- (1) $\text{Min}(A) = \bigcap_{\alpha \in A} \text{Min}(\alpha)$ is a closed, convex, Γ -invariant, quasi-dense subspace of X that splits as a product $Y \times Z$, where Z is isometric to $\mathbb{R}^n$,
- (2) G acts geometrically on Y by projection of the action of Γ on $Y \times Z$, and
- (3) A acts geometrically on Z by restriction of the action of Γ on $Y \times Z$ (moreover, Z is the convex hull of some orbit Ax_0 of X).

Here a subset X' of a metric space X is said to be *quasi-dense* if there exists a number $N > 0$ such that each point of X is N -close to some point of X' , i.e., $B(X', N) = X$.

Using Theorem 1.1 and results above, we prove Theorem 1.2 on structure of $CAT(0)$ groups.

Proof of Theorem 1.2. Let Γ be a $CAT(0)$ group which acts geometrically on a proper $CAT(0)$ space X .

Set $\Gamma_0 = \Gamma$.

Suppose that Γ_0 has infinite center. Then the center of Γ_0 is an abelian subgroup of the $CAT(0)$ group Γ_0 and finitely generated by Lemma 2.5. Hence there exists a central element $\gamma_1 \in \Gamma_0$ with the order $o(\gamma_1) = \infty$. Then the isometry γ_1 of X is semi-simple, since every element of the $CAT(0)$ group Γ_0 is a semi-simple isometry of X by Lemma 2.4. Also γ_1 is a hyperbolic isometry, because if γ_1 is an elliptic isometry then γ has a bounded orbit by Lemma 2.3, which contradicts that the action of Γ_0 on X is proper, since $o(\gamma_1) = \infty$.

Now Γ_0 is finitely generated by Lemma 2.2 and γ_1 is a central element of Γ_0 and a hyperbolic isometry of X . Hence by Theorem 1.1, there exist a normal subgroup $\Gamma_1 \subset \Gamma_0$, an element $\delta_1 \in \Gamma_0$ and a number $k_1 \in \mathbb{N}$ such that

- (i₁) $\Gamma_0 = \Gamma_1 \rtimes \langle \delta_1 \rangle$,
- (ii₁) $\Gamma_1 \rtimes \langle \delta_1^{k_1} \rangle = \Gamma_1 \times \langle \gamma_1 \rangle$ is a finite-index subgroup of Γ_0 and
- (iii₁) Γ_0/Γ_1 is isomorphic to $\mathbb{Z}$.

Here $\Gamma_1 \times \langle \gamma_1 \rangle$ acts geometrically on the CAT(0) space X , since $\Gamma_1 \times \langle \gamma_1 \rangle$ is a finite-index subgroup of Γ_0 by (ii₁). Hence by Theorem 2.6, $\text{Min}(\gamma_1)$ is a closed convex Γ -invariant quasi-dense subspace of X that splits as a product $X_1 \times Z_1$, where Z_1 is isometric to $\mathbb{R}$, and Γ_1 acts geometrically on X_1 by projection of the action of Γ_0 on $X_1 \times Z_1$. Here we note that X_1 is a CAT(0) space, because X_1 is a convex subspace of the CAT(0) space X (cf. [2] and [3]). Hence Γ_1 acts geometrically on the CAT(0) space X_1 and Γ_1 is a CAT(0) group.

Suppose that Γ_1 has infinite center. Then by the same argument as above, there exists a central element $\gamma_2 \in \Gamma_1$ with $o(\gamma_2) = \infty$, which is a hyperbolic isometry of X_1 . Here the CAT(0) group Γ_1 is finitely generated. Hence by Theorem 1.1, there exist a normal subgroup $\Gamma_2 \subset \Gamma_1$, an element $\delta_2 \in \Gamma_1$ and a number $k_2 \in \mathbb{N}$ such that

- (i₂) $\Gamma_1 = \Gamma_2 \rtimes \langle \delta_2 \rangle$,
- (ii₂) $\Gamma_2 \rtimes \langle \delta_2^{k_2} \rangle = \Gamma_2 \times \langle \gamma_2 \rangle$ is a finite-index subgroup of Γ_1 and
- (iii₂) Γ_1/Γ_2 is isomorphic to $\mathbb{Z}$.

Then

$$\begin{aligned}\Gamma &= \Gamma_0 = \Gamma_1 \rtimes \langle \delta_1 \rangle \\ &= (\Gamma_2 \rtimes \langle \delta_2 \rangle) \rtimes \langle \delta_1 \rangle.\end{aligned}$$

Also

$$\begin{aligned}\Gamma &= \Gamma_1 \rtimes \langle \delta_1 \rangle \supset \Gamma_1 \rtimes \langle \delta_1^{k_1} \rangle \\ &= \Gamma_1 \times \langle \gamma_1 \rangle = (\Gamma_2 \rtimes \langle \delta_2 \rangle) \times \langle \gamma_1 \rangle \\ &\supset (\Gamma_2 \rtimes \langle \delta_2^{k_2} \rangle) \times \langle \gamma_1 \rangle \\ &= (\Gamma_2 \times \langle \gamma_2 \rangle) \times \langle \gamma_1 \rangle \\ &= \Gamma_2 \times \langle \gamma_1 \rangle \times \langle \gamma_2 \rangle.\end{aligned}$$

Thus $\Gamma_2 \times A_2$ is a finite-index subgroup of Γ , where $A_2 = \langle \gamma_1 \rangle \times \langle \gamma_2 \rangle$ which is isomorphic to $\mathbb{Z}^2$.

Also Γ_2 is a CAT(0) group which acts geometrically on some CAT(0) space X_2 by Theorem 2.6.

By iterating this argument, we obtain subgroups $\Gamma = \Gamma_0 \supset \Gamma_1 \supset \cdots \supset \Gamma_n$, elements $\delta_{i+1}, \gamma_{i+1} \in \Gamma_i$ and $k_{i+1} \in \mathbb{N}$ for $i = 0, \dots, n-1$ such that

- (1) γ_{i+1} is an element of the center of Γ_i with $o(\gamma_{i+1}) = \infty$ for $i = 0, \dots, n-1$,
- (2) $\Gamma_i = \Gamma_{i+1} \rtimes \langle \delta_{i+1} \rangle$ for $i = 0, \dots, n-1$,
- (3) $\Gamma_{i+1} \rtimes \langle \delta_{i+1}^{k_{i+1}} \rangle = \Gamma_{i+1} \times \langle \gamma_{i+1} \rangle$ is a finite-index subgroup of Γ_i ,

- (4) Γ_i/Γ_{i+1} is isomorphic to $\mathbb{Z}$ for $i = 0, \dots, n-1$,
- (5) $\Gamma = (\dots(((\Gamma_n \rtimes \langle \delta_n \rangle) \rtimes \langle \delta_{n-1} \rangle) \rtimes \langle \delta_{n-2} \rangle) \dots) \rtimes \langle \delta_1 \rangle$,
- (7) $\Gamma_n \times A_n$ is a finite-index subgroup of Γ where $A_n = \langle \gamma_1 \rangle \times \dots \times \langle \gamma_n \rangle$ which is isomorphic to $\mathbb{Z}^n$.

Here this process must terminate, i.e., Γ_n has finite center for some number n , because $A_n \cong \mathbb{Z}^n$ is an abelian subgroup of the CAT(0) group Γ and every abelian subgroup of a CAT(0) group is finitely generated by Lemma 2.5. $\square$

3. EXAMPLE

We introduce an easy example of a CAT(0) group.

Example. Let $\Gamma = \langle a, b \mid ab^2 = b^2a \rangle$ and let $X = \mathbb{R}^2$ the euclidean plane. We consider the action of the group Γ on X defined by

$$\begin{aligned} a \cdot (x, y) &= (x, y + 1) \\ b \cdot (x, y) &= (x + 1, -y) \end{aligned}$$

for any $(x, y) \in \mathbb{R}^2 = X$. Then $D = [0, 1] \times [-\frac{1}{2}, \frac{1}{2}] \subset \mathbb{R}^2$ is a fundamental domain, $\Gamma D = X$ and Γ acts geometrically on X . Here we note that X/Γ is a Klein bottle and the group Γ is a CAT(0) group which is the fundamental group of the Klein bottle.

Then $\gamma_0 := b^2$ is a center of the CAT(0) group Γ and a hyperbolic isometry of X . Here we obtain that

- (i) $\Gamma = \langle a \rangle \rtimes \langle b \rangle$,
- (ii) $\langle a \rangle \rtimes \langle b^2 \rangle = \langle a \rangle \times \langle b^2 \rangle$ is a finite-index subgroup of Γ which is isomorphic to $\mathbb{Z}^2$ and
- (iii) $\Gamma/\langle a \rangle$ is isomorphic to $\mathbb{Z}$.

4. REMARKS ON CAT(0) GROUPS

We introduce some remarks on CAT(0) groups.

Let Γ be a CAT(0) group. By Theorem 1.2, Γ has a structure as

$$\Gamma = (\dots(((\Gamma_n \rtimes \langle \delta_n \rangle) \rtimes \langle \delta_{n-1} \rangle) \rtimes \langle \delta_{n-2} \rangle) \dots) \rtimes \langle \delta_1 \rangle,$$

where Γ_n is a CAT(0) group with finite-center, and Γ contains a finite-index subgroup $\Gamma_n \times A$ where A is isomorphic to $\mathbb{Z}^n$.

Here we first note that the CAT(0) group Γ_n may contain a finite-index CAT(0) subgroup with infinite-center in general. For example, all Coxeter groups are CAT(0) groups with finite-center and affine Coxeter groups contain a finite-index CAT(0) subgroup with infinite-center which is isomorphic to $\mathbb{Z}^n$ for some n . Also in general, the CAT(0) group Γ may contain a finite-index subgroup $\Gamma' \times A'$ where A' is isomorphic to $\mathbb{Z}^{n'}$ and $n' \geq n$.

If the CAT(0) group Γ splits as $\Gamma = \Gamma_1 \times \cdots \times \Gamma_m$, then each factor Γ_i is also a CAT(0) group (cf. [14, Theorem 6]) and each factor Γ_i has a structure as

$$\Gamma_i = (\cdots (((\Gamma_{i,n_i} \rtimes \langle \delta_{i,n_i} \rangle) \rtimes \langle \delta_{i,n_i-1} \rangle) \rtimes \langle \delta_{i,n_i-2} \rangle) \cdots) \rtimes \langle \delta_{i,1} \rangle,$$

where Γ_{i,n_i} is a CAT(0) group with finite-center, and Γ_i contains a finite-index subgroup $\Gamma_{i,n_i} \times A_i$ where A_i is isomorphic to $\mathbb{Z}^{n_i}$ for each $i = 1, \dots, m$. Here $\Gamma = \Gamma_1 \times \cdots \times \Gamma_m$ contain a finite-index subgroup

$$(\Gamma_{1,n_1} \times \cdots \times \Gamma_{m,n_m}) \times (A_1 \times \cdots \times A_m).$$

We note that $\Gamma_{1,n_1} \times \cdots \times \Gamma_{m,n_m}$ has finite center and $A_1 \times \cdots \times A_m$ is isomorphic to $\mathbb{Z}^{n_1 + \cdots + n_m}$.

Also as the argument in [13, p.909], there exists a finite-index subgroup $G_1 \times \cdots \times G_l$ of the CAT(0) group Γ such that each factor G_j is infinite and each factor G_j does not contain a finite-index subgroup $G_{j1} \times G_{j2}$ where G_{j1} and G_{j2} are infinite. Here the decomposition process terminates and l is finite (cf. [13, p.909]). Then we note that each factor G_j is isomorphic to $\mathbb{Z}$ or has finite center. Let t be the number as G_j has finite center for $1 \leq j \leq t$ and G_j is isomorphic to $\mathbb{Z}$ for $t+1 \leq j \leq l$, and let $n'' = l - t$. Then $G_1 \times \cdots \times G_t$ has finite center, $A'' = G_{t+1} \times \cdots \times G_l$ is isomorphic to $\mathbb{Z}^{n''}$ and $(G_1 \times \cdots \times G_t) \times A''$ is a finite-index subgroup of Γ .

Here we can use the Flat Torus Theorem and splitting theorems (cf. [3], [14] and [16]) in each case above.

For example, if the CAT(0) group Γ acts geometrically on a CAT(0) space X , then the finite-index subgroup $(G_1 \times \cdots \times G_t) \times A''$ also acts geometrically on X and the Flat Torus Theorem and splitting theorems implies that X contains a quasi-dense subspace X' which splits as a product $(X_1 \times \cdots \times X_t) \times Z$ such that there exists a geometric action of G_j on X_j for each $j = 1, \dots, t$ and Z is isometric to $\mathbb{R}^{n''}$ (cf. [14]). Then the boundary ∂X of the CAT(0) space X (cf. [3], [5], [17] and [19]) is homeomorphic to the spherical join $(\partial X_1 * \cdots * \partial X_t) * \partial Z$, where ∂Z is homeomorphic to an $(n'' - 1)$ -dimensional sphere.

By these arguments, CAT(0) groups with finite-center that does not contain a finite-index subgroup which splits as a product of infinite groups are essential and important for research on CAT(0) spaces on which a CAT(0) group acts geometrically and their boundaries.

Also, Theorem 1.2 (5) on structure of CAT(0) groups implies that CAT(0) groups with finite-center are essential and important for research on CAT(0) groups.

REFERENCES

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV-VI, Masson, Paris, 1981.
- [2] P. Bowers and K. Ruane, *Boundaries of nonpositively curved groups of the form $G \times \mathbb{Z}^n$* , Glasgow Math. J. 38 (1996), 177–189.
- [3] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [4] L. S. Charlap, *Bieberbach groups and flat manifolds*, Universitext. Springer-Verlag, New York, 1986.
- [5] C. B. Croke and B. Kleiner, *Spaces with nonpositive curvature and their ideal boundaries*, Topology 39 (2000), 549–556.
- [6] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293–324.
- [7] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.
- [8] M. W. Davis and T. Januszkiewicz, *Hyperbolization of polyhedra*, J. Diff. Geom. **34** (1991), 347–388.
- [9] E. Ghys and P. de la Harpe (ed), *Sur les groupes hyperboliques d’après Mikhael Gromov*, Progr. Math. vol. 83, Birkhäuser, Boston MA, 1990.
- [10] M. Gromov, *Hyperbolic groups*, Essays in group theory (S. M. Gersten, ed.), M.S.R.I. Publ. 8, 1987, pp. 75–264.
- [11] M. Gromov, *Asymptotic invariants for infinite groups*, Geometric Group Theory (G.A. Niblo and M.A. Roller, eds.), LMS Lecture Notes, vol. 182, Cambridge University Press, Cambridge, 1993, pp. 1–295.
- [12] P. de la Harpe, *Topics in geometric group theory*, Chicago Lectures in Math. Ser., The University of Chicago Press.
- [13] T. Hosaka, *Minimality of the boundary of a right-angled Coxeter system*, Proc. Amer. Math. Soc., 137 (2009), 899–910.
- [14] T. Hosaka, *On splitting theorems for $CAT(0)$ spaces and compact geodesic spaces of non-positive curvature*, preprint, arXiv:math.GR/0405551v1 (2004).
- [15] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [16] N. Monod, *Superrigidity for irreducible lattices and geometric splitting*, J. Amer. Math. Soc. 19 (2006), 781–814.
- [17] C. Mooney, *Examples of non-rigid $CAT(0)$ groups from the category of knot groups*, preprint, arXiv:0706.1581v1 (2007).
- [18] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, The Ohio State University, 1988.
- [19] J. M. Wilson, *A $CAT(0)$ group with uncountably many distinct boundaries*, J. Group Theory 8 (2005), 229–238.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, UTSUNOMIYA
UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN
E-mail address: hosaka@cc.utsunomiya-u.ac.jp